



Fermi National Accelerator Laboratory

FERMILAB-TM-2059

Passage Through a Transverse Magnetic Slab

Leo Michelotti

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

October 1998

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Distribution

Approved for public release; further dissemination unlimited.

Copyright Notification

This manuscript has been authored by Universities Research Association, Inc. under contract No. DE-AC02-76CHO3000 with the U.S. Department of Energy. The United States Government and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government Purposes.

Passage through a transverse magnetic slab

Leo Michelotti

July 29, 1998

The purpose of this short note is to derive the simple but remarkable result that the transverse impulse given to a particle passing through a “slab” of homogeneous, stationary, transverse magnetic field depends only on the properties of the slab and the particle’s charge, and not at all on its initial state or its mass. Let a constant magnetic field, \vec{B} , exist in a region bounded by two parallel planes separated by a distance, l . If the triple, $(\hat{e}_1, \hat{e}_2, \hat{e}_3)$, is an orthonormal frame of vectors with \hat{e}_3 orthogonal to the planes, then the condition that \vec{B} is “transverse” can be written as either $\vec{B} \cdot \hat{e}_3 = 0$ or $\vec{B} = B_1 \hat{e}_1 + B_2 \hat{e}_2$. We want to show that the transverse impulse, $(\hat{e}_1 \hat{e}_1 + \hat{e}_2 \hat{e}_2) \cdot \Delta \vec{p}$, given to a particle passing through the region is independent of its dynamical coordinates at the point of entry. (Since the field is homogeneous, this is obviously true with regard to its position.)

Begin with the Lorentz equation.

$$\dot{\vec{p}} = e \vec{v} \times \vec{B} ,$$

where e is the particle’s electric charge. The impulse given to the particle is obtained by integrating this through the field region.

$$\begin{aligned} \Delta \vec{p} &= e \int dt (\vec{v} \times \vec{B}) \\ &= e \int d\vec{s} \times \vec{B} \\ &= e \left(\int d\vec{s} \right) \times \vec{B} \\ &= e \vec{L} \times \vec{B} , \end{aligned} \tag{1}$$

where \vec{L} is a displacement vector connecting the entry point to the exit point. Because $\vec{B} \cdot \hat{e}_3 = 0$, we can introduce the auxiliary field \vec{U} , defined by

$$\begin{aligned} \vec{B} &= \hat{e}_3 \times \vec{U}, \text{ or, equivalently} \\ \vec{U} &= \vec{B} \times \hat{e}_3 . \end{aligned}$$

Note in passing that

$$\text{if } \vec{B} = B_1 \hat{e}_1 + B_2 \hat{e}_2, \text{ then } \vec{U} = B_2 \hat{e}_1 - B_1 \hat{e}_2 .$$

That is, $U_1 = B_2$ and $U_2 = -B_1$. We rewrite Eq.(1) in terms of \vec{U} .

$$\begin{aligned}\vec{L} \times \vec{B} &= \vec{L} \times (\hat{e}_3 \times \vec{U}) \\ &= (\vec{L} \cdot \vec{U})\hat{e}_3 - (\vec{L} \cdot \hat{e}_3)\vec{U}\end{aligned}\quad (2)$$

It will be convenient to make the angular dependence explicit and express everything in terms of l , the length of the field region. We write $\vec{L} = L\hat{n}$ and use $l = \vec{L} \cdot \hat{e}_3 = L\hat{n} \cdot \hat{e}_3$. Substituting these and Eq.(2) back into Eq.(1), we obtain.

$$\Delta\vec{p} = (el) \left[\left(\hat{n} \cdot \vec{U} / \hat{n} \cdot \hat{e}_3 \right) \hat{e}_3 - \vec{U} \right]$$

Only the longitudinal component, Δp_3 , depends on orientation.

$$\begin{aligned}\Delta p_3 &= el \frac{\hat{n} \cdot \vec{U}}{\hat{n} \cdot \hat{e}_3} \\ &= el \frac{\hat{n} \cdot (\vec{B} \times \hat{e}_3)}{\hat{n} \cdot \hat{e}_3}\end{aligned}$$

However, in practice, we never compute this component directly but merely infer it from the transverse components and the fact that magnetic fields cannot alter the energy of a particle. The transverse components are free from any angular dependence.

$$\begin{aligned}\Delta p_1 &= -elU_1 \\ &= -eB_2l \\ \Delta p_2 &= -elU_2 \\ &= eB_1l\end{aligned}$$

These equations express the desired result, as their right hand sides depend only on properties of the field region and the particle's charge. Usually, we divide by a reference momentum,

$$p_{\text{ref}} = e|B\rho|_{\text{ref}}$$

and rewrite them as follows.

$$\begin{aligned}\Delta p_1/p_{\text{ref}} &= -B_2l/|B\rho|_{\text{ref}} \\ \Delta p_2/p_{\text{ref}} &= B_1l/|B\rho|_{\text{ref}}\end{aligned}$$

This is an exact result, but it is extended via the ‘‘thin element’’ approximation, in which we indulge ourselves with the polite fiction that, if l is sufficiently small, the particle's transverse coordinates remain ‘‘essentially constant’’ in passing through the field. The approximation enables us to allow transverse variations in B_2 and B_1 ,

$$\begin{aligned}\Delta p_1/p_{\text{ref}} &\approx -B_2(x, y)l/|B\rho|_{\text{ref}} , \\ \Delta p_2/p_{\text{ref}} &\approx B_1(x, y)l/|B\rho|_{\text{ref}} .\end{aligned}$$

These equations then form the basis of a number of ‘‘kick codes’’ for tracking particles.